



Ozark Center
COMPREHENSIVE BEHAVIORAL HEALTH SERVICES
An entity of Freeman Health System

The Facts about Buprenorphine for Treatment of Opioid Addiction



How do you escape from the misery and risks of drug addiction? Most people cannot do it on their own. They need help to return to normal, healthy living. One important form of help is called medication-assisted treatment.

This form of treatment has three equally important components:

- Medication
- Counseling
- Support from family and friends

Buprenorphine is one type of medication used for treatment of opioid addiction. Your doctor or substance abuse treatment provider may have recommended it for you, or you may already take it. This booklet will give you the facts you need to know.

NOTE: Important words often used in treatment are introduced in this booklet in bold type.

Opioid addiction is a disease.

Opioids are drugs that slow down the actions of the body, such as breathing and heartbeat. Getting their name from opium, a drug made from the poppy plant, opioids also affect the brain to increase pleasant feelings.

Opioid medications are prescribed to treat pain and other health problems such as severe coughing. Heroin is an illegal opioid that is smoked, snorted or injected to get a good feeling, often called a “rush” or “high.” People also sometimes seek this feeling by taking large doses of prescription opioids.

If you take opioids, you can become **tolerant** to them. This means that more of the drug is needed to obtain its effects. It is also possible to become **dependent** on opioids, meaning if you stop taking them, you will feel sick. This sickness is called **withdrawal**. Dependence is not the same as **addiction**, but sometimes dependence leads to addiction. The signs of addiction are:

- **Craving** – The mind develops an overwhelming desire for the drug.
- **Loss of control** – It becomes harder to say no to using the drug. Use is compulsive and continues even though the drug causes harm.

Tolerance and dependence are common side effects of prescribed opioid medication and sometimes can be managed under a doctor’s care. Tolerance and dependence also are side effects of misuse of opioids. Addiction is not likely to develop in a person using medication properly, but it sometimes happens. Addiction usually occurs through misuse. Some people are at higher risk of addiction because of their genes, temperament or personal situation.

“I’d been shot on the streets. I’d been in detox. It was jails, institutions... Death, I knew, was imminent for me... so I started my recovery process... It’s been 15 years since then, and my life has never been better.”

Walston B.

Not everyone who uses opioids gets addicted. However, after you have become addicted, it is difficult to stop using opioids because the cravings are strong and the fear of withdrawal is great.

Opioid addiction can be treated.

Opioid addiction is a **chronic disease**, like heart disease or diabetes, meaning it is a medical condition for life. It cannot be cured, but it can be managed. A person with addiction can regain a healthy, productive life. If you are like most people, though, you cannot walk away from addiction on your own. **Treatment** – the care of medical professionals and substance use treatment providers – **can help**.

Treatment helps you give up the problem drug. It helps you get through withdrawal and cope with cravings. Treatment also helps you change addictive thinking into nonaddictive, healthful patterns. It can help you move away from other harmful behaviors, too, such as drinking alcohol or abusing other drugs besides the problem opioid.

Just as importantly, treatment helps you address life issues that may be tied to your addiction, such as feelings of low self-worth, a bad situation at work or home, or spending time with people who use drugs. In short, treatment helps you move into a healthy, addiction-free lifestyle – into a way of living referred to as **recovery**.



Treatment may include medication.

Medication-assisted treatment includes the use of medication along with counseling and other support. Treatment that includes medication is often the best choice for opioid addiction.

If you are addicted, medication allows you to regain a normal state of mind, free of drug-induced highs and lows. It frees you from thinking about the drug all the time. It can reduce problems of withdrawal and craving. These changes can give you the chance you need to focus on the lifestyle changes that lead back to healthy living.

Taking medication for opioid addiction is like taking medication to control heart disease or diabetes. It is NOT the same as substituting one addictive drug for another. Used properly, the medication does NOT create a new addiction. It helps you manage your addiction so that the benefits of recovery can be maintained.

Buprenorphine is a medication used to treat opioid addiction.

Buprenorphine is one of three medications commonly used to treat opioid addiction. The other two are **methadone** and **naltrexone**. Cost varies for the different medications. You may need to take this into account when considering your treatment options.

The person who takes buprenorphine feels normal, not high. However, the brain thinks it is receiving the problem opioid, so withdrawal symptoms don't appear. Buprenorphine also reduces cravings. If cravings continue to be a problem, your doctor will adjust your medication or help you find other ways to reduce them.

You take buprenorphine as a pill that dissolves under the tongue. You do NOT chew or swallow it. There are two forms. **Suboxone**® contains buprenorphine plus another medication called naloxone. The naloxone is added to prevent abuse – it brings on withdrawal in people who abuse buprenorphine by injecting it. **Subutex**® contains only buprenorphine. This form is prescribed if you should not take naloxone for any reason, such as if you are allergic to it or are pregnant.

The pill is taken once a day. Over time, the dose interval may stay at once a day or change to every other day.

The main advantages of buprenorphine are:

- You are unlikely to overdose on buprenorphine if you take it properly.
- Buprenorphine is long-acting. This means that after an initial period, your doctor may have you take the pill every other day rather than once a day.
- Doctors can prescribe buprenorphine so that you can take doses at home. Important: Not all doctors have approval to prescribe this medication, and not all doctors provide counseling for addiction. Daily check-in at a treatment center can be also helpful in recovery, making a treatment center is the best place for some people to receive medication for opioid addiction.

Buprenorphine may be prescribed if it is the right choice for you. _____

Before you are given the first dose of any medication for opioid addiction, your doctor will ask you questions about your addiction, health and other problems. You will get a drug test – usually a check of urine or saliva. You also will have a physical exam and tests for diseases that are common to people who have been abusing drugs. Your liver will be checked to make sure the medication can be safely taken. If buprenorphine is safe and appropriate for you, your doctor may recommend it.

You and your doctor or substance use treatment provider will decide together on a treatment plan. The plan describes the medication routine, the counseling and other services that will be provided, and the rules that must be followed. These rules will be explained to you, and you will get them in writing.

Buprenorphine must be used carefully. _____

Your doctor will prescribe a low dose to start taking after withdrawal symptoms begin. Your doctor may increase dose levels as needed. Always take doses exactly as prescribed by your doctor.

Buprenorphine can make you feel drowsy at first. You should not drive or perform other high-risk tasks until you know how this medication affects you. If drowsiness continues to be a problem, your doctor may adjust dose levels.

You may take buprenorphine for days, months or years – as long as it is needed to prevent relapse. However, you should be checked often by a doctor if you have liver disease.

If you are stable in recovery and want to stop taking buprenorphine, you must do so slowly, over time. This is called **tapering**. Tapering works best with the help of your doctor or substance use treatment provider, after progress has been made in treatment.

WARNINGS

- While taking this medication, you should NOT take other medications without consulting your doctor first.
- While taking this medication, you should NOT use illegal drugs, drink alcohol or take sedatives, tranquilizers or other drugs that slow breathing. Taking any of these substances in large amounts along with buprenorphine can lead to overdose or death.
- Buprenorphine kept at home must be locked in a safe place to prevent accidental use by others, especially children.
- If you are a woman and are pregnant or breastfeeding, methadone is safer than buprenorphine for mother and child. In special circumstances, doctors may recommend the naloxone-free form of buprenorphine instead of methadone. Ask your doctor for more information.
- Liver problems are rare but can occur. If you have a pre-existing liver condition, your doctor should conduct regular tests on your liver.

In some people, buprenorphine has **side effects**, or feelings of discomfort or sickness from taking the medicine. Buprenorphine's side effects are usually mild and go away after the medication is taken for a while.

If you have side effects, you should NOT stop taking the medication. Instead, talk with your doctor or substance use treatment provider. An adjustment in dosage or a change in medication may help. There are simple ways you can reduce some of the side effects, too.

Common side effects and simple ways to reduce them.

Body aches, headaches and cold- or flu-like symptoms. Check with your doctor about over-the-counter medicines you may take.

Dizziness. Stand up slowly. Call your doctor if problems persist.

Constipation. Drink more water and juice. Eat food with fiber. Exercise more.

Sweating. Shower often. Dress in layers.

Sleep problems, including fatigue. Take the pill in the morning. Avoid naps. Go to bed at the same time every night. Exercise. Do not drink caffeine after lunchtime.

Upset stomach or vomiting. Take the pill after you have eaten. Take an antacid product as directed by your doctor.

Mood swings. Exercise more. Do fun things that do not involve the old drug lifestyle. Relax. Talk to your substance use treatment provider.

Serious side effects. For extreme stomach pain, vomiting or diarrhea, contact your doctor right away. Also seek help if the following side effects appear, because they may indicate serious liver problems:

- Dark or tea-colored urine
- Bad stomachache
- Light-colored bowel movements
- Yellowing in the whites of the eyes
- Yellow skin

Counseling can help. _____

Medication is one part of treatment for opioid addiction.

For many people, another important part is **counseling** – the opportunity to talk with a professional, either one-on-one or in a group with others in treatment.



Through counseling, you learn about the motivations and behaviors that led to your opioid addiction. You learn how to commit to a more healthful lifestyle. You gain support and skills while working with others to manage your recovery long term.

Counseling can provide you with encouragement and motivation to stick to treatment. It can help you learn how to make healthy decisions, handle setbacks and stress, and move forward with your life.

In **group counseling**, you connect with others in treatment and make new friends who don't use drugs. You can also get these benefits from **support groups**, or informal meetings of people facing similar challenges.

Family and friends are important too. _____

It is very hard to go through recovery alone. Love and encouragement from family and friends can help you make the decision to enter treatment and stick with it.

Some treatment programs offer counseling for your loved ones because your addiction may have caused pain and anger or feelings of shame and hopelessness.

Counseling is a useful way for family and friends to learn more about your situation, how to help and how to handle the problems your addiction has caused them. It is a safe place for them to express feelings and to find out what help is available for everyone affected.

There are support groups for family and friends, too. These are safe places to share information and encourage others who have loved ones who are dealing with addiction.

Many people with an opioid addiction regain normal, healthy lives. The combination of medication, counseling and support can help you, your loved ones and your friends.

ADDICTION

What's true and what's not.

Addiction is a disease. It cannot be cured, but it can be treated with medication, counseling and support from family and friends. Addiction is NOT a sign of weakness. It is NOT TRUE that all you need to do to kick addiction is "be strong."

The goal of medication-assisted treatment is to recover from addiction. It does NOT replace one addictive drug with another. It provides a safe, controlled level of medication to overcome the use of a problem opioid.

A substance use treatment provider must obtain your informed consent (agreement in writing) before sharing information about you with others. However, there are two exceptions to this privacy rule: (1) if it appears that you may harm yourself or others and (2) if you have been ordered into treatment by the courts. To learn more about your privacy rights, talk to your substance use treatment provider.

Recovery is possible. It takes work, though. After treatment is finished, everything is NOT automatically fine again. Recovery takes commitment every day, through treatment and beyond.

BUPRENORPHINE

What's true and what's not.

Buprenorphine helps you think and function normally. It is legal and taken under a doctor's care. It is NOT just another drug to abuse.

Buprenorphine is produced under safe conditions and sold legally. There is no risk of getting tainted doses, which can happen with street drugs. Also, risk of overdose on this medication is very small.

You can stop taking buprenorphine when you are ready. You can become dependent on buprenorphine, as with many medications taken over time. For this reason, if you wish to stop taking buprenorphine, you should work with a substance use treatment provider to taper off. This prevents withdrawal symptoms from appearing.

"The treatment center I was at had a lot of education about addiction. They talked about it as a disease and not a moral failing... I had a medical condition... It was a big relief for me to hear that and to be able to realize that recovery was possible."

Tom C.

Support groups and information.

- This is not a complete list. Listing here does not mean Ozark Center or the Substance Abuse and Mental Health Services Administration (SAMHSA) endorse any of the organizations.
- Some support groups have abstinence-only policies and do not look favorably on medication-assisted treatment. The programs listed here do not have such policies, but individual group meetings vary. You may need to try several support groups to find the right one.
- Some support programs are just for people with a substance use disorder, and others allow families and friends to attend meetings or have separate meetings for them. Check with each organization for details.
- An internet-based support group may be your best option if no groups meet in your community. Another option is to contact Alcoholics Anonymous (aa.org) to find out whether AA meetings in your community are open to people in recovery from substances other than alcohol.

Dual Recovery Anonymous
draonline.org

Rational Recovery
rational.org or 530.621.4374

Women for Sobriety, Inc.
womenforsobriety.org or
215.536.8026

LifeRing
lifering.org or 800.811.4142

Secular Organizations for Sobriety
sossobriety.org or 323.666.4295

**Buprenorphine physicians and
treatment program locator**
800.662.HELP (4357) (English and
Español) 800.487.4889 TDD (for
hearing impaired)
findtreatment.samhsa.gov

**National Alliance of Advocates for
Buprenorphine Treatment**
naabt.org

SMART Recovery
smartrecovery.org or 866.951.5357

Free booklets.

These booklets are available at samhsa.gov. Ask your doctor or substance use treatment provider for more information.

- The Facts About Naltrexone for Treatment of Opioid Addiction (SMA) 09-4444 (also in Spanish)
- Medication-Assisted Treatment of Opioid Use Disorder Pocket Guide (SMA16-4892PG)
- Decisions in Recovery: Treatment for Opioid Use Disorders (SMA16-4993)
- Faces of Change: An Illustrated Booklet for Consumers (SMA08-4174)
- What Is Substance Abuse Treatment? A Booklet for Families (SMA08-4126) Spanish version: (SMA08-4098)

“What I’ve found in my life is that the more people I tell that I’m a person in recovery..., then the more people will be able to carry that message to their friends and family and say, ‘Hey, I know somebody who doesn’t use drugs anymore who used to, and his life has gotten better. Maybe you can give this a try.’”

Dan P.



Acknowledgements

Information in this document was originally prepared for the Substance Abuse and Mental Health Services Administration (SAMHSA) by the Knowledge Application Program, a joint venture of the CDM Group, Inc., and JBS International, Inc., under contract number 270-04-7049 with SAMHSA, U.S. Department of Health and Human Services (HHS).

ATENCIÓN: si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística. Llame al 417-347-7600 (TTY: 1- 800-735-2966).
注意: 如果您使用繁體中文, 您可以免費獲得語言援助服務。請致電 417-347-7600 (TTY: 1- 800-735-2966).